

EDITORIAL

Factors for advancing emergency airway management practice

Hui-Chun Ku¹, Shih-Yi Lee^{2,3,*}

¹Department of Life Science, Fu Jen Catholic University, 242062 Taipei, Taiwan

²Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, MacKay Memorial Hospital, 10499 Taipei, Taiwan

³MacKay Junior College of Medicine, Nursing, and Management, 11260 Taipei, Taiwan

***Correspondence**

leesy15538@yahoo.com.tw
(Shih-Yi Lee)

Abstract

Compiling factors from scientific evidence of airway management in the emergency setting is a strategy for improving patient outcomes. Review and original articles in this special issue highlight the essential elements that deserve special attention in clinical practice, including airway assessments before artificial airway establishment, rapid sequence induction, continuous monitoring of vital signs after airway intubation, and management of artificial airway-associated discomfort.

Keywords

Capnography; Difficult airway; Difficult mask ventilation; Difficult intubation; Emergency airway management; Rapid sequence intubation; Tube-related sore throat

Emergency airway management is a crucial aspect in critical care. Relieving ventilation insufficiency and airway obstruction saves lives; airway management strategies must also be optimized to improve airway management quality [1–3]. Factors for advancing emergency airway management practice include airway assessments before artificial airway establishment [4, 5], rapid sequence induction [6], use of device to verify the position of the artificial airway and to monitor ventilation and pulmonary circulation after intubation [7, 8], and management of artificial airway-induced discomfort [9].

Difficult airway (DA) resulting in difficult mask ventilation or difficult intubation complicates emergency airway management. The difficulty in establishing sufficient oxygenation through bag-valve-mask ventilation potentially leads to complications such as eye and eyelid injury, airway trauma, nerve injury, gastric insufflation by overinflation (inflation pressure: >20 cmH₂O), vomiting and aspiration of gastric contents into the airway [10, 11]; multiple attempts to establish artificial airways possibly lead to adverse events such as airway trauma, esophageal intubation, aspiration, hypoxemia, hypotension, dysrhythmia, and cardiac arrest [12, 13].

A quick review among patients requiring airway management is a strategy to deduce a DA [14]. Successfully identifying physical features suggestive of a DA is crucial for accurate preparation before artificial airway intubation and for planning a safe emergency airway management [15]. The emergency airway assessment tool should be quick and effective. Age-related changes in the structure and function surrounding the airway ultimately result in the development of anatomic features of DA in elderly people [16]. One original study in this special issue evaluated whether age-related changes in the head and neck anatomy among apparently normal adult patients lead to difficulty in emergency airway management; hence, rapid decisions related to preparation for emergency airway

management may be guided by patient age [17].

In addition to airway obstruction in apparently normal patients, pathologic airway obstruction can limit airway flow [18–20]. Head and neck disease may change the internal anatomy surrounding supra- and infra-glottic airways [21, 22]. Vascular anomalies in the area below the larynx may cause airway narrowing and are not visible during examination with a laryngoscope during trachea intubation [23]. Therefore, multidisciplinary preoperative discussion is required to plan for special therapeutic considerations before, during, and after surgical reconstruction for vascular-related airway obstruction; moreover, physicians should remember to anticipate the signs of unexpected airway obstruction and prepare the equipment to resolve the problem when it occurs [24]. Furthermore, the next article in the special issue reviewed airway obstruction in deep neck infection. Deep neck infection potentially causes airway narrowing from the mouth along the trachea to the carina because of trismus, laryngeal edema, and the mass effect caused by the abscess in the suprahyoid space, infrahyoid space, or that spreading from the neck to the mediastinum [25]. Securing the airway in patients undergoing surgical intervention for deep neck infection is challenging because they usually exhibit limited mouth opening, airway distortion, tissue edema, or cervical immobility. Assessing the risk of a potential DA and planning the most appropriate airway management method are critical in managing patients with deep neck infection.

Rapid sequence intubation is the cornerstone of emergency airway management [5]. Administering a sedative agent immediately followed by a neuromuscular blocking agent to produce rapid unconsciousness and paralysis provides optimum intubation conditions while minimizing aspiration. Successful and safe tracheal intubations with anesthesia induction should prevent morbidity and serious consequences of a failure in airway establishment. However, upper airway obstruction eas-

ily occurs in critically ill patients with reduced consciousness levels [26]. Anesthesia induction in an unconscious patient can further result in loss of the pharyngeal tone and falling-back tongue, leading to airway narrowing and in turn desaturation, which complicates the tracheal intubation procedure, especially in the case of a DA [27, 28]. Moreover, physicians must be aware of the hemodynamic effects of general anesthesia agents used for critically ill patients during rapid sequence intubation [29]. An ideal sedative agent is effective, safe, and rapid acting; acts in a short duration; and allows optimum intubation conditions [30]. Because of its influence on catecholamine secretion and sympathetic nervous system stimulation, ketamine has been presumed to be a beneficial alternative for hemodynamically unstable patients [31]. The article by Bakhsh *et al.* [32] in this special issue discussed a study comparing the change in hemodynamic parameters between two commonly available induction agents for rapid sequence intubation, namely etomidate and ketamine [32].

Carbon dioxide is a crucial physiological indicator of ventilation, pulmonary circulation, and aerobic metabolism [33]. Among the methods for confirming endotracheal tube placement (*i.e.*, auscultation of the chest and epigastrium, visualization of symmetrical thoracic movement and fogging flow, ultrasonography, and radiography), capnography that detects end-tidal carbon dioxide in exhaled breath is the gold standard [34]. The readings of capnography are influenced by ventilation and cardiac output levels [33]; hence, simply using the values of end-tidal carbon dioxide detected by capnography is not considered very reliable for verifying the position of an endotracheal tube during emergency airway management [35, 36]. Real-time portable quantitative waveform capnography is another vital method in patients receiving emergency airway management and requiring continuous monitoring of ventilation and pulmonary circulation in near-arrest, during resuscitation, and during transport to critical care units after return of spontaneous circulation [5, 37, 38]. The review article by Huang *et al.* [39] in this special issue pointed out the components required for interpreting quantitative capnography waveform for physicians to deliver prompt accurate treatment in response to the clinical changes in patients requiring emergency airway management [39].

Physicians should also pay attention to artificial airway-induced discomfort because intubated patients are unable to voice their needs [40–42]. The care of postoperative sore throat has been extensively studied [9, 43]; however, the medical care of discomfort from prolonged endotracheal intubation in critical patients has not been extensively explored [44–47]. The final article in this special issue compared the therapeutic effects of two common pain relievers, oral acetaminophen and local lidocaine application, on endotracheal tube-related sore throat in critically ill patients [48].

This special issue is intended to give the readers of *Signa Vitae-Journal of Anesthesia, Intensive Care, Emergency and Pain Medicine* a quick review of the factors required to obtain favorable clinical outcomes in patients requiring emergency airway management. The special issue also attempts to provide a perspective for future studies involved in emergency airway management. The introduction of new technologies, such as video laryngoscopy with a hyperangulated blade and laryngeal

mask airway, for airway management promotes evolutionary changes in the DA definition, with changes to target study groups depending on the equipment and environment used for practicing emergency airway management [49–56]. Moreover, studies focusing on an intervention for tracheal tube-related sore throat, a major source of stress but an easily neglected discomfort in critically ill patients [57, 58], are warranted to fill the unmet need.

AUTHOR CONTRIBUTIONS

SYL and HCK designed the study, SYL and HCK searched, reviewed, and analyzed the articles. HCK drafted the manuscript. SYL and HCK completed and proofread the article.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

ACKNOWLEDGMENT

The authors acknowledge the institutional support from the Ministry of Science and Technology. We also would like to express our gratitude to all those who helped us during building up the special issue, and the entire peer reviewers for their opinions and suggestions.

FUNDING

This research was funded by the Ministry of Science and Technology, Taiwan, grant number MOST 109-2320-B-030-006-MY3.

CONFLICT OF INTEREST

The authors declare no conflict of interest. Shih-Yi Lee is serving as one of the Editorial Board members of this journal. We declare that Shih-Yi Lee had no involvement in the peer review of this article and has no access to information regarding its peer review. Full responsibility for the editorial process for this article was delegated to PB.

REFERENCES

- [1] Jaber S, Amraoui J, Lefrant JY, Arich C, Cohendy R, Landreau L, *et al.* Clinical practice and risk factors for immediate complications of endotracheal intubation in the intensive care unit: a prospective, multiple-center study. *Critical Care Medicine*. 2006; 34: 2355–2361.
- [2] Jaber S, Jung B, Corne P, Sebbane M, Muller L, Chanques G, *et al.* An intervention to decrease complications related to endotracheal intubation in the intensive care unit: a prospective, multiple-center study. *Intensive Care Medicine*. 2010; 36: 248–255.
- [3] Perbet S, De Jong A, Delmas J, Futier E, Pereira B, Jaber S, *et al.* Incidence of and risk factors for severe cardiovascular collapse after endotracheal intubation in the ICU: a multicenter observational study. *Critical Care*. 2015; 19: 257.
- [4] Lee SY, Chien DK, Huang MY, Huang CH, Shih SC, Wu KM, *et al.* Patient-specific factors associated with difficult mask ventilation in the

- emergency department. *International Journal of Gerontology*. 2017; 11: 263–266.
- [5] Apfelbaum JL, Hagberg CA, Connis RT, Abdelmalak BB, Agarkar M, Dutton RP, *et al.* 2022 American society of anesthesiologists practice guidelines for management of the difficult airway. *Anesthesiology*. 2022; 136: 31–81.
- [6] Okubo M, Gibo K, Hagiwara Y, Nakayama Y, Hasegawa K, Japanese Emergency Medicine Network Investigators. The effectiveness of rapid sequence intubation (RSI) versus non-RSI in emergency department: an analysis of multicenter prospective observational study. *International Journal of Emergency Medicine*. 2017; 10: 1.
- [7] Jooste R, Roberts F, Mndolo S, Mabedi D, Chikumbanje S, Whitaker DK, *et al.* Global capnography project (GCAP): implementation of capnography in Malawi—an international anaesthesia quality improvement project. *Anaesthesia*. 2019; 74: 158–166.
- [8] Gelb AW, Morriss WW, Johnson W, Merry AF, International Standards for a Safe Practice of Anesthesia Workgroup. World health organization-world federation of societies of anaesthesiologists (WHO-WFSA) international standards for a safe practice of anesthesia. *The Canadian Journal of Anesthesia*. 2018; 65: 698–708.
- [9] AlQahtani RM, Abdalla M, Azzam YH, Elsherif AA, Altulayh RI. Pharmacological interventions for post-operative sore throat (POST): a network meta-analysis. *Signa Vitae*. 2021; 17: 169–177.
- [10] Lee SY, Kuo CY, Wei DH, Ku HC, Chung RJ, Chang WH. Perspective view for mask design. *Health Technol*. 2017; 1: 3.
- [11] Cook TM, MacDougall-Davis SR. Complications and failure of airway management. *British Journal of Anaesthesia*. 2012; 109: i68–i85.
- [12] Hasegawa K, Shigemitsu K, Hagiwara Y, Chiba T, Watase H, Brown CA, *et al.* Association between repeated intubation attempts and adverse events in emergency departments: an analysis of a multicenter prospective observational study. *Annals of Emergency Medicine*. 2012; 60: 749–754.e2.
- [13] Cook TM, Woodall N, Harper J, Benger J, Fourth National Audit Project. Major complications of airway management in the UK: results of the fourth national audit project of the royal college of anaesthetists and the difficult airway society. Part 2: intensive care and emergency departments. *British Journal of Anaesthesia*. 2011; 106: 632–642.
- [14] Apfelbaum JL, Hagberg CA, Caplan RA, Blitt CD, Connis RT, Nickinovich DG, *et al.* Practice guidelines for management of the difficult airway: an updated report by the American society of anesthesiologists task force on management of the difficult airway. *Anesthesiology*. 2013; 118: 251–270.
- [15] Roth D, Pace NL, Lee A, Hovhannisyan K, Warenits AM, Arrich J, *et al.* Airway physical examination tests for detection of difficult airway management in apparently normal adult patients. *Cochrane Database of Systematic Reviews*. 2018; 5: CD008874.
- [16] Lee SY, Shih SC, Leu YS, Chang WH, Lin HC, Ku HC. Implications of age-related changes in anatomy for geriatric-focused difficult airways. *International Journal of Gerontology*. 2017; 11: 130–133.
- [17] Liao EC, Chang WH, Yu CH, Chau YP, Sun FJ, Lai WJ, *et al.* Predictors of difficult endotracheal intubation in the emergency department: a single-center pilot study. *Signa Vitae*. 2021; 17: 77–84.
- [18] Drašković B, Uram Benka A, Drašković D, Marić D, Simić D. Anaesthesiological problems in children with congenital laryngeal stenosis. *Signa Vitae*. 2010; 5: 38–39.
- [19] Kralik S, Škarić I, Butković D, Mikecin L, Kondza K, Jakobović J. Serious complications of an obstructive upper airway infection in a young child. *Signa Vitae*. 2009; 4: 30–32.
- [20] Yamakawa K, Dohgomor H, Furusawa T. Early morning upper airway discomfort and appearance on two x ray films. *Signa Vitae*. 2007; 2: 21–22.
- [21] Lee J, Naing K, Yeo ZZ, Chong PH. The use of continuous positive airway pressure ventilation in the palliative management of stridor in a head and neck cancer patient. *Journal of Pain and Symptom Management*. 2019; 58: e3–e5.
- [22] Sesterhenn AM, Iwinska-Zelder J, Dalchow CV, Bien S, Werner JA. Acute haemorrhage in patients with advanced head and neck cancer: value of endovascular therapy as palliative treatment option. *The Journal of Laryngology & Otology*. 2006; 120: 117–124.
- [23] Škarić I, Jakobović J, ILIĆ MK, Kondza K, Babić I. Variation of vascular ring as a cause of extubation failure. *Signa Vitae*. 2008; 3: 29–31.
- [24] Lee SY, Chao CL, Hung ST, Lim HK, Ku HC. Airway management in vascular central airway obstruction: a literature review. *Signa Vitae*. 2021; 17: 8–17.
- [25] Lim HK, Wang JM, Hung ST, Ku HC. A dangerous cause of airway obstruction—deep neck infection. *Signa Vitae*. 2021; 17: 4–9.
- [26] McPherson K, Stephens RC. Managing airway obstruction. *British Journal of Hospital Medicine*. 2012; 73: C156–C160.
- [27] Law JA, Broemling N, Cooper RM, Drolet P, Duggan LV, Griesdale DE, *et al.* The difficult airway with recommendations for management—part 2—the anticipated difficult airway. *The Canadian Journal of Anesthesia*. 2013; 60: 1119–1138.
- [28] Law JA, Broemling N, Cooper RM, Drolet P, Duggan LV, Griesdale DE, *et al.* The difficult airway with recommendations for management—part 1—difficult tracheal intubation encountered in an unconscious/induced patient. *The Canadian Journal of Anesthesia*. 2013; 60: 1089–1118.
- [29] Lah K, Miljenko K, Grmed S. Rapid sequence intubation in the pre-hospital setting—difference between trauma and nontrauma patients. *Signa Vitae*. 2010; 5: 34–39.
- [30] Hampton JP. Rapid-sequence intubation and the role of the emergency department pharmacist. *American Journal of Health-System Pharmacy*. 2011; 68: 1320–1330.
- [31] Shin TG, Jo S, Hwang SY, Jo II, Lee TR, Yoon H, *et al.* Ketamine use for endotracheal intubation in severe sepsis and septic shock. *Signa Vitae*. 2018; 14: 24–30.
- [32] Bakhsh A, Alnashri M, Alawami F, Aseel R, Almaghthawi M, Alrahaili G, *et al.* Changes in hemodynamic parameters with the use of etomidate versus ketamine induction in the emergency department. *Signa Vitae*. 2021; 17: 85–92.
- [33] Zorica NT. Clinical applications of capnography. *Signa Vitae*. 2008; 3: S44–S45.
- [34] Gökhan I, Süleyman D, Sedat T, Ömer B, Hüseyin S, Sezai O. Validation of tracheal intubation of wire-reinforced endotracheal tube with ultrasonography. *Signa Vitae*. 2018; 14: 20–23.
- [35] Li J. Capnography alone is imperfect for endotracheal tube placement confirmation during emergency intubation. *The Journal of Emergency Medicine*. 2001; 20: 223–229.
- [36] Vaghadia H, Jenkins LC, Ford RW. Comparison of end-tidal carbon dioxide, oxygen saturation and clinical signs for the detection of oesophageal intubation. *Canadian Journal of Anaesthesia*. 1989; 36: 560–564.
- [37] Hatlestad D. Capnography as a predictor of the return of spontaneous circulation. *Emergency medical services*. 2004; 33: 75–80.
- [38] Long B, Koefman A, Vivirito MA. Capnography in the emergency department: a review of uses, waveforms, and limitations. *The Journal of Emergency Medicine*. 2017; 53: 829–842.
- [39] Huang CH, Wei KH. Applications of capnography in airway management outside the operating room. *Signa Vitae*. 2021; 17: 18–24.
- [40] Grap MJ, Blecha T, Munro C. A description of patients’ report of endotracheal tube discomfort. *Intensive and Critical Care Nursing*. 2002; 18: 244–249.
- [41] Rotondi AJ, Chelluri L, Sirio C, Mendelsohn A, Schulz R, Belle S, *et al.* Patients’ recollections of stressful experiences while receiving prolonged mechanical ventilation in an intensive care unit. *Critical Care Medicine*. 2002; 30: 746–752.
- [42] Topolovec-Vranic J, Canzian S, Innis J, Pollmann-Mudryj MA, McFarlan AW, Baker AJ. Patient satisfaction and documentation of pain assessments and management after implementing the adult nonverbal pain scale. *American Journal of Critical Care*. 2010; 19: 345–354.
- [43] Ahmed A, Abbasi S, Ghafoor HB, Ishaq M. Postoperative sore throat after elective surgical procedures. *Journal of Ayub Medical College*. 2007; 19: 12–14.
- [44] Samuelson KA. Adult intensive care patients’ perception of endotracheal tube-related discomforts: a prospective evaluation. *Heart & Lung*. 2011; 40: 49–55.
- [45] Rahu MA, Grap MJ, Cohn JF, Munro CL, Lyon DE, Sessler CN. Facial expression as an indicator of pain in critically ill intubated adults during endotracheal suctioning. *American Journal of Critical Care*. 2013; 22: 412–422.
- [46] Shinn JR, Kimura KS, Campbell BR, Sun Lowery A, Wootten CT,

- Garrett CG, *et al.* Incidence and outcomes of acute laryngeal injury after prolonged mechanical ventilation. *Critical Care Medicine*. 2019; 47: 1699–1706.
- [47] Lee SY, Wu CL, Kuo LK, Lai CT, Hsu CP, Hwung HY, *et al.* The effects of xylocaine spray for pain control caused by endotracheal tube in critical care. *International Journal of Gerontology*. 2012; 6: 11–15.
- [48] Lim HK, Lee SY, Wu CW, Lai JCY, Ho YH, Ku HC. Comparative study between oral acetaminophen and lidocaine spray on endotracheal tube-related sore throat in adult intensive care. *Signa Vitae* 2021; 17: 71–76.
- [49] Heidegger T. Management of the difficult airway. *New England Journal of Medicine*. 2021; 384: 1836–1847.
- [50] Cooper RM. Preparation for and management of “failed” laryngoscopy and/or intubation. *Anesthesiology*. 2019; 130: 833–849.
- [51] Ramachandran SK, Mathis MR, Tremper KK, Shanks AM, Kheterpal S. Predictors and clinical outcomes from failed laryngeal mask airway unique™: a study of 15,795 patients. *Anesthesiology*. 2012; 116: 1217–1226.
- [52] Lee DK, Shih DH, Park SM, Kim YH, Park SO, Lee YH. I-gel as a first-line airway device in the emergency room for patients with out-of-hospital cardiac arrest. *Signa Vitae*. 2018; 14: 61–65.
- [53] Lin SC, Hsu SC, Weng YM, Kuo CI, Cheng CW, Kuo CW. Dose pre-hospital laryngeal mask airway use has a survival benefit in non-shockable cardiac arrest? *Signa Vitae*. 2014; 9: 27–32.
- [54] Wass TC, Jacob AK, Kopp SL, Torsher LC. A prospective randomized high fidelity simulation center based side-by-side comparison analyzing the success and ease of conventional versus new generation video laryngoscope technology by inexperienced laryngoscopists. *Signa Vitae*. 2011; 6: 36–45.
- [55] Chien LC, Hsu HC, Lin CH, Cheng CF, Tung YC, Hung HC, *et al.* Use of an intubating laryngeal mask airway on out-of-hospital cardiac arrest patients in a developing emergency medical service system. *Journal of the Formosan Medical Association*. 2012; 111: 24–29.
- [56] Liu YC, Huang WC, Tan EC, Huang SS, Wang YK, Chu YC. Practice and outcomes of airway management in patients with cervical orthoses. *Journal of the Formosan Medical Association*. 2022; 121: 108–116.
- [57] Lee SY, Hung CL, Lee JH, Shih SC, Weng YL, Chang WH, *et al.* Attaining good end-of-life care in intensive care units in Taiwan—the dilemma and the strategy. *International Journal of Gerontology*. 2009; 3: 26–30.
- [58] Gillis J, Lai CT, Lin ML, Lee SY. Connecting palliative care and intensive care in Taiwan. *International Journal of Gerontology*. 2008; 2: 33–34.

How to cite this article: Hui-Chun Ku, Shih-Yi Lee. Factors for advancing emergency airway management practice. *Signa Vitae*. 2022; 18(6): 1-4. doi:10.22514/sv.2022.032.