

ORIGINAL RESEARCH



Admission rates and outcomes of elderly patients in emergency department observation units. A Spanish multicentre study

Francisco Javier Montero-Pérez^{1,*}, Inmaculada Bajo-Fernández¹, Juan González del Castillo², Guillermo Burillo-Putze³, Cesáreo Fernández², Sira Aguiló⁴, Aitor Alquézar-Arbé⁵, Javier Jacob⁶, Pere Llorens^{7,8}, Osvaldo Jorge Troiano Ungerer⁹, Patricia Eiroa-Hernández³, Patricia Parra-Esquivel¹⁰, Mabel Coromoto Suárez Pineda¹¹, Martín Ruiz Grinspan¹², María Florencia Poblete Palacios⁴, Matilde González Tejera¹³, María Bóveda García¹⁴, Mónica Veguillas Benito⁷, Ferran Llopis⁶, Rocío Muñoz Martos¹⁵, Valle Toro Gallardo¹⁶, Rodrigo Javier Gil Hernández¹⁷, Jorge Pedraza García¹⁸, Carmen Lucena Aguilera¹, María Dolores Pulfer¹⁹, María Eugenia Rodríguez²⁰, Marta Iglesias Vela²¹, María Rodríguez Romero²², Alba Morant Gimenez²³, Òscar Miró⁴,
On behalf of the researchers of the SIESTA network

¹Emergency Department, Reina Sofía University Hospital of Córdoba, Maimonides Biomedical Research Institute of Córdoba (IMIBIC), 14004 Córdoba, Spain

²Emergency Department, Hospital Clínico San Carlos, IDISSC, Complutense University, 28040 Madrid, Spain

³Emergency Department, University Hospital of the Canary Islands, University of La Laguna, 38320 La Laguna, Tenerife, Canary Islands, Spain

⁴Emergency Department, Hospital Clínic, IDIBAPS, University of Barcelona, 08036 Barcelona, Spain

⁵Emergency Department, Hospital de la Santa Creu i Sant Pau, 08041 Barcelona, Spain

⁶Emergency Department, Hospital Bellvitge University Hospital, L'Hospitalet de Llobregat, 08907 Barcelona, Spain

⁷Emergency Department, Alicante Biomedical and Health Research Institute (ISABIAL), Dr. Balmis University General Hospital, 03010 Alicante, Spain

⁸Clinic Medicine Department, Miguel Hernández University, 03202 Elche, Spain

⁹Emergency Department, Hospital Santa Tecla, 43003 Tarragona, Spain

¹⁰Emergency Department, North Tenerife Hospital, 38430 Icod de los Vinos, Tenerife, Canary Islands, Spain

¹¹Emergency Department, Reina Sofía University General Hospital of Murcia, 30003 Murcia, Spain

¹²Emergency Department, University Hospital "del Henares", 28822 Madrid, Spain

¹³Emergency Department, Elche University General Hospital, 03203 Elche, Spain

¹⁴La Fe University and Polytechnic Hospital, 46026 Valencia, Spain

¹⁵Emergency Department, Hospital de la Axarquía, 29700 Vélez-Málaga, Spain

¹⁶Emergency Department, Hospital Regional de Málaga, 29010 Málaga, Spain

¹⁷Emergency Department, Hospital Santa Bárbara, 42002 Soria, Spain

¹⁸Emergency Department, Hospital Valle de los Pedroches, 14400 Pozoblanco, Córdoba, Spain

¹⁹University General Hospital Gregorio Marañón, 28007 Madrid, Spain

²⁰University Hospital of Burgos, 09001 Burgos, Spain

²¹University Care Complex of León, 24071 León, Spain

²²Hospital General Universitario Morales Meseguer, 30008 Murcia, Spain

²³Hospital Francisc de Borja, 46701 Gandía, Spain

***Correspondence:** franciscoj.montero.sspa@juntadeandalucia.es (Francisco Javier Montero-Pérez)

Abstract

Background: Emergency Department Observation Units (EDOUs) provide short-term treatment and monitoring for patients who require further evaluation. EDOUs may help reduce unnecessary hospital admissions in elderly adults, but their selection criteria and impact on outcomes remain unclear. The study, thus, aimed to identify clinical factors associated with using EDOUs in patients aged ≥ 65 years and to evaluate the relationship between EDU care and short-term clinical outcomes. **Methods:** We analysed data from the Emergency Department and Elder Needs (EDEN) cohort, which included all emergency department visits by patients aged ≥ 65 years across 48 Spanish hospitals during seven days. We divided patients into two groups: those managed in an EDU (EDU group) and those managed without observation care (non-EDU group). We examined demographic and clinical characteristics, emergency diagnoses, and 30-day outcomes. Multivariable logistic regression identified factors independently associated with EDU use and subsequent outcomes. **Results:** Among 23,955 visits, 6393 (26.7%) involved EDU management. Patients in the EDU group were more likely to be ≥ 80 years and to present with tachypnoea, bradycardia, a Glasgow Coma Scale score < 15 , and anaemia. These variables showed significant independent associations with EDU care. Compared to the non-EDU group, the EDU group had higher rates of hospital admission (adjusted odds Ratio (aOR) = 2.4; 95% confidence interval (CI): 2.0–2.8) and 30-day readmission (aOR = 1.6; 95% CI: 1.2–2.3), but similar rates of 30-day ED revisit, prolonged hospital stay, and mortality. Patient selection varied across centres and often lacked standardized protocols. **Conclusions:** EDU care for elderly adults in Spanish emergency departments typically involves patients with greater clinical complexity. Although associated with higher hospital admission and readmission rates, EDU use did not correlate with worse short-term outcomes. These findings support the need for standardized EDU admission criteria in geriatric populations.

Keywords

Emergency department; Observation unit; Older adults; Geriatric emergency care; Hospital admission; Short-term outcomes

1. Introduction

As the global population ages, older adults account for a growing proportion of emergency department (ED) visits [1, 2]. In many countries, patients aged ≥ 65 represent up to 30% of ED activity [3]. These patients often present with higher risk and complexity due to functional impairment, cognitive decline, polypharmacy, and multiple chronic conditions [4, 5].

Emergency Department Observation Units (EDOUs) offer short-term clinical observation and management for patients who require further assessment before a disposition decision. These units benefit patients whose condition may evolve or need additional testing or response monitoring. EDOUs typically operate with stays under 48 hours [4] and provide an alternative to inpatient admission when discharge is not immediately safe or feasible.

EDOUs may benefit older patients by avoiding unnecessary hospitalization while offering close observation. Geriatric conditions such as falls, delirium, syncope, heart failure, or respiratory infections often fall within the scope of EDU care [4–11]. Evidence suggests that EDOUs can improve patient flow, reduce costs, and maintain safety, particularly when combined with protocolized management [12–18].

Despite these potential advantages, few studies have examined how ED teams use EDOUs for older patients or how this decision affects outcomes [19–21]. Most prior research has focused on general adult populations or EDU performance metrics rather than patient selection.

This study, therefore, aims to identify clinical and functional factors associated with the decision to manage patients aged

≥ 65 years in an EDU. We also assess how EDU use relates to key short-term outcomes, including hospitalization, ED revisits, and mortality. Our findings may support future development of evidence-based criteria for EDU admission in geriatric emergency care.

2. Methods

2.1 Description of the EDEN challenge and SIESTA network

The Emergency Department and Elder Needs (EDEN) challenge originated from the Spanish Investigators on Emergency Situation TeAm (SIESTA) research network [3, 4], which includes 52 emergency departments (EDs), approximately 20% of all public EDs in Spain. Its main objective is to increase knowledge about the sociodemographic, organizational, baseline, clinical, care-related, and evolutionary aspects of patients aged 65 and older who attend Spanish EDs.

To achieve this, we created a multipurpose registry that included all patients aged ≥ 65 who visited participating EDs between 01 April and 07 April 2019 (seven consecutive days), regardless of the reason for consultation. This registry constitutes the EDEN cohort. Extended patient recruitment and follow-up details have been published elsewhere [22–24].

2.2 EDEN-20 study design

The present study, EDEN-20, is a secondary analysis of patients in the EDEN cohort. For this analysis, we included data from the 48 EDs with an operational Emergency Department

Observation Unit (EDOU) at the time of patient inclusion. Patients were classified into two groups: those admitted to an observation unit (EDOU group) and those who were not (non-EDOU group). Both groups included patients discharged from the ED or admitted to the hospital. The classification was based solely on whether observation unit care was used at any point during the ED stay.

In addition, we collected information about the structural and functional characteristics of each participating EDOU. These included hospital type (primary/secondary vs. tertiary), number of available observation beds, organizational structure (open, closed, or mixed models), and functional classification (Types I–IV). Type I units follow standardized protocols and are managed directly by ED staff; Type II units apply partial protocols or are jointly managed; Types III and IV represent more complex or less standardized observation models.

We analysed patients' sociodemographic variables, baseline functional and cognitive status, and polypharmacy. These included nine variables: age, sex, comorbidities, walking ability, cognitive impairment, mode of arrival to the ED, referral source, and presence of polypharmacy.

We considered the following comorbid conditions: arterial hypertension, dyslipidaemia, uncomplicated or complicated (organ failure) diabetes mellitus, chronic lung disease, heart failure, ischemic heart disease, chronic kidney disease, stroke, dementia, cancer with or without metastasis, peripheral vascular disease, connective tissue disease, venous thromboembolism, human immunodeficiency virus (HIV)/acquired immunodeficiency syndrome (AIDS), alcoholism, leukaemia or lymphoma, and moderate or severe chronic liver disease.

We also collected eight variables related to physical examination: hypotension, hypertension, significant tachypnoea, tachycardia, significant bradycardia, hypoxemia, fever, and Glasgow Coma Scale (GCS) score <15. In addition, we analysed 14 laboratory abnormalities: leucocytosis, leukopenia, erythrocytosis, anaemia, thrombocytosis, thrombocytopenia, hypoglycaemia, elevated serum creatinine, hyponatremia, hyponatremia, hyperkalaemia, hypokalaemia, and elevated serum lactate. We dichotomized laboratory results using common clinical cut-off points.

Given the patient acuity and presentation heterogeneity, laboratory and diagnostic testing were performed only when clinically indicated. Initial ED vital signs were similarly dichotomized using standard clinical thresholds. Fever was defined as temperature $\geq 38^\circ\text{C}$, tachycardia as heart rate >120 bpm, bradycardia as <50 bpm, tachypnoea as respiratory rate >20 breaths per minute, hypoxemia as peripheral capillary oxygen saturation (SpO_2) $<90\%$, hypotension as systolic blood pressure (BP) <90 mmHg, and hypertension as systolic BP ≥ 160 mmHg.

Additionally, we analysed the most frequent primary emergency diagnoses in both study groups. These Emergency diagnoses were registered using ICD-10 codes (*10th revision of the International Statistical Classification of Diseases and Related Health Problems*). The length of stay in the ED was also registered, and was computed as the period between registration in the emergency department and the decision to discharge home or hospital admission for the non-EDOU group. In the case of the EDOU group, the time spent in this

unit was also considered in the calculation.

2.3 Outcomes

We examined five short-term outcomes: (1) hospital admission following ED or EDOU management; (2) prolonged hospital stay, defined as inpatient length of stay ≥ 7 days; (3) 30-day ED revisit (only for discharged patients); (4) 30-day hospital readmission; and (5) 30-day all-cause mortality. All outcomes were obtained from electronic health records via standardized data collection protocols.

2.4 Statistical analysis

We conducted a descriptive analysis comparing the EDOU and non-EDOU groups. Categorical variables are reported as absolute frequencies and percentages, and continuous variables as medians with interquartile ranges (IQR). We used chi-square or Fisher's exact tests to compare categorical variables, and Mann-Whitney U tests for continuous variables.

We developed two multivariable logistic regression models. The first identified independent factors associated with EDOU admission. The second assessed the association between EDOU care and each of the short-term outcomes. We included in the models all variables showing significant differences in univariate analysis and those deemed clinically relevant. We report unadjusted odds ratios (OR) and adjusted OR (aOR) with respective 95% confidence intervals (CI), and consider p -values < 0.05 statistically significant. All analyses were performed using IBM SPSS Statistics version 17.0 (IBM, Armonk, NY, USA).

3. Results

Table 1 presents the structural and functional characteristics of the participating EDOUs, providing context for their utilization.

The EDEN-20 study included 23,955 ED visits across 48 Spanish hospitals. Of these, 6393 visits (26.7%) involved patients admitted to an EDOU (EDOU group), while 17,562 (73.3%) did not (non-EDOU group) (Fig. 1). The median age of all patients was 78 years (IQR: 13); 10,433 patients (43.6%) were aged 80 or older, and 55% were women.

The median ED length of stay was 8.3 hours (IQR: 15.22) for the EDOU group and 2.7 hours (IQR: 3.24) for the non-EDOU group. A total of 1078 patients (17%) in the EDOU group had ED stays longer than 24 hours (including time in the EDOU), compared to only 226 patients (1.3%) in the non-EDOU group (OR = 16.0; 95% CI: 14.0–18.0; $p < 0.001$).

Compared to non-EDOU patients, those in the EDOU group were older (median age 81 vs. 77 years; $p < 0.001$), with a higher proportion aged ≥ 80 years, a greater burden of comorbidity, worse functional capacity (as measured by the Barthel Index), more frequent mobility impairment, and higher rates of cognitive decline. Patients in the EDOU group more often arrived by ambulance and were more frequently referred by a physician or from another hospital. Table 2 shows these variables along with their unadjusted ORs and statistical significance.

The same table also presents data on vital signs. The

TABLE 1. Characteristics of participating hospitals and observation units in the EDEN-20 study.

Variables	Primary/Secondary Hospitals (n = 26)	Tertiary Hospitals (n = 22)	Total (N = 48)
Number of observation beds (median (IQR))	17 (15)	38 (27)	24 (27)
Organizational structure (n (%))			
Open model	18 (69.2%)	11 (50.0%)	29 (60.4%)
Closed model	4 (15.4%)	5 (22.7%)	9 (18.8%)
Mixed model	4 (15.4%)	6 (27.3%)	10 (20.8%)
Functional classification (n (%))			
Type I	12 (46.2%)	14 (63.6%)	26 (54.2%)
Type II	12 (46.2%)	7 (31.8%)	19 (39.6%)
Type III	1 (3.8%)	1 (4.6%)	2 (4.2%)
Type IV	1 (3.8%)	0 (0.0%)	1 (2.1%)

IQR: interquartile ranges.




FIGURE 1. Flowchart for patient inclusion in the EDEN-20 cohort. EDOU: Emergency Department Observation Unit; EDEN: Emergency Department and Elder Needs. Note: bold text is used to highlight key cohort definitions and main outcome categories.

TABLE 2. Baseline demographics, functional and clinical characteristics among patients admitted and not admitted to Emergency Department Observation Units.

Variables	EDOU N = 6393 n (%)	Non-EDOU N = 17,562 n (%)	OR (95% CI)	p-value
Sociodemographic characteristics				
Age group (yr)				
<80 (reference)	2888 (45.2)	10,634 (60.6)	-	<0.001
≥80	3505 (54.8)	6928 (39.4)	2.0 (1.8–2.0)	
Sex (female) ^a	3141 (50.3)	9785 (56.9)	0.8 (0.7–0.8)	<0.001
Arrival to ED				<0.001
Their own (reference)	3653 (57.0)	14,206 (80.9)	3.2 (3.0–3.4)	<0.001
Non-medicalized or medicalized ambulance	2740 (43.0)	3356 (19.1)		
Referred to the ED				
Own initiative of the patient or caregiver (reference)	3823 (60.0)	12,932 (74.0)	2.0 (1.8–2.0)	<0.001
Referred from primary care or by a medical specialist (other than primary care) or another hospital	2570 (40.2)	4630 (26.0)		
Baseline status				
Comorbidity presence	6128 (96.0)	15,794 (90.0)	2.6 (2.3–3.0)	<0.001
Functional capacity (by Barthel Index)				<0.001
Independent (100 points) (reference)	3444 (54.0)	12,550 (71.5)	-	<0.001
Mild or moderate (60–95 points)	1974 (31.0)	3680 (21.0)	2.0 (1.8–2.1)	
Severe or complete (<60 points)	975 (15.0)	1332 (7.6)	2.4 (2.4–2.9)	
Walking ability				<0.001
Alone with no help (reference)	3744 (58.6)	13,249 (75.4)	-	<0.001
Need help	1983 (31.0)	3444 (19.6)	2.7 (2.4–3.0)	
Unable to walk	666 (10.4)	869 (4.9)	2.0 (1.9–2.2)	
Baseline cognitive decline	1253 (19.6)	1968 (11.2)	2.0 (1.8–2.1)	<0.001
Polypharmacy (≥5 drugs)	4737 (74.1)	10,419 (59.3)	2.0 (1.8–2.1)	<0.001
Vitals at ED arrival				
Arterial hypertension (systolic arterial pressure >160 mmHg) ^b	1053 (18.3)	2251 (23.0)	0.8 (0.7–0.8)	<0.001
Arterial hypotension (systolic arterial pressure <90 mmHg) ^b	174 (3.0)	133 (1.3)	2.3 (1.8–2.9)	<0.001
Fever (≥38 °C) ^c	102 (4.5)	68 (1.6)	2.9 (2.1–3.9)	<0.001
Significant tachypnoea (>20 breaths per minute) ^d	504 (16.4)	259 (4.7)	4.0 (3.4–4.7)	<0.001
Tachycardia (>120 beats per minute) ^e	303 (5.5)	226 (2.4)	2.4 (2.0–2.9)	<0.001
Significant bradycardia (<50 beats per minute) ^e	112 (2.0)	137 (1.4)	1.4 (1.1–1.8)	<0.001
Hypoxemia (peripheral arterial oxygen saturation ≤90%) ^f	692 (14.1)	456 (6.2)	2.5 (2.2–2.8)	<0.001
Glasgow coma scale score <15 ^g	285 (8.2)	191 (2.3)	3.8 (3.2–4.6)	<0.001

TABLE 2. Continued.

Variables	EDOU N = 6393 n (%)	Non-EDOU N = 17,562 n (%)	OR (95% CI)	p-value
Analytical data				
Leucocytosis (>11,000/microL) ^h	1831 (31.3)	1770 (22.5)	1.6 (1.4–1.7)	<0.001
Leukopenia (<4000/microL) ^h	167 (3.0)	231 (3.0)	1.0 (0.8–1.2)	0.800
Erythrocytosis (hemoglobin ≥16.5 g/dL in men; ≥16 g/dL in women) ⁱ	162 (2.6)	183 (1.1)	2.5 (2.0–3.0)	<0.001
Anaemia (hemoglobin <9 g/dL) ⁱ	364 (7.5)	225 (4.0)	2.0 (1.7–2.4)	<0.001
Thrombocytosis (>400,000/microL) ^j	309 (5.3)	310 (4.0)	1.4 (1.2–1.6)	<0.001
Thrombocytopenia (<100,000/microL) ⁱ	200 (3.4)	261 (3.3)	1.0 (0.9–1.2)	0.738
Hyperglycaemia (≥180 mg/dL) ^k	1031 (18.1)	1049 (13.4)	1.4 (1.3–1.6)	<0.001
Hypoglycaemia (<70 mg/dL) ^k	61 (1.1)	84 (1.1)	1.0 (0.7–1.4)	0.100
Elevated serum creatinine (>1.4 mg/dL) ^k	1529 (27.0)	1444 (19.0)	1.6 (1.5–1.7)	<0.001
Hypernatremia (>145 mEq/L) ^l	205 (3.5)	169 (2.2)	1.7 (1.3–2.0)	<0.001
Hyponatremia (<135 mEq/L) ^l	913 (15.8)	960 (12.4)	1.3 (1.2–1.5)	<0.001
Hyperkalaemia (>5.5 mEq/L) ^m	219 (4.0)	193 (2.6)	1.6 (1.3–1.9)	<0.001
Hypokalaemia (<3.5 mEq/L) ^m	410 (7.3)	404 (5.4)	1.4 (1.2–1.6)	<0.001
Serum lactate (>2 mmol/L or >18 mg/dL) ⁿ	468 (30.0)	365 (29.3)	1.0 (0.9–1.2)	0.900

EDOU: Emergency Department Observation Unit; OR: unadjusted odds ratio (for EDOU admission); CI: confidence intervals; ED: Emergency Department.

p-values in bold denote statistical significance ($p < 0.05$).

^avariable registered in 6249 of the EDOU group and 17,1393 episodes of the No-EDOU group.

^bvariable registered in 5757 of the EDOU group and 9878 episodes of the No-EDOU group.

^cvariable registered in 2286 of the EDOU group and 4239 episodes of the No-EDOU group.

^dvariable registered in 3068 of the EDOU group and 5541 episodes of the No-EDOU group.

^evariable registered in 5531 of the EDOU group 9616 episodes of the No-EDOU group.

^fvariable registered in 4919 of the EDOU group and 7405 episodes of the No-EDOU group.

^gvariable registered in 3467 of the EDOU group and 8339 episodes of the No-EDOU group.

^hleukocyte count was requested or registered in 5856 of the EDOU group and 7878 episodes of the No-EDOU group.

ⁱhemoglobin determination was requested or registered in 4824 of the EDOU group and 5770 episodes of the No-EDOU group.

^jplatelet count was requested or registered in 5799 of the EDOU group and 7827 episodes of the No-EDOU group.

^kplasmatic glucose and serum creatinine were requested or registered in 5686 of the EDOU group and 7808 episodes of the No-EDOU group.

^lplasmatic sodium was requested or registered in 5785 of the EDOU group and 7768 episodes of the No-EDOU group.

^mplasmatic potassium was requested or registered in 5611 of the EDOU group and 7442 episodes of the No-EDOU group.

ⁿserum lactate was requested or registered in 1581 of the EDOU group and 1245 episodes of the No-EDOU group.

presence of arterial hypotension, arterial hypertension, significant tachypnoea, tachycardia, bradycardia, hypoxemia, and a Glasgow Coma Scale (GCS) score <15 was significantly associated with EDOU admission. The strongest associations were observed for tachypnoea (OR = 4.0; 95% CI: 3.4–4.7) and GCS <15 (OR = 3.8; 95% CI: 3.2–4.6).

Regarding laboratory abnormalities (Table 2), erythrocytosis (OR = 2.5; 95% CI: 2.0–3.0) and anaemia (OR = 2.0; 95% CI: 1.7–2.4) showed the strongest associations with EDOU admission. Of the 31 variables analysed in the EDEN-20 dataset, 27 showed statistically significant differences between the EDOU and non-EDOU groups.

Table 3 displays the 10 most common ED diagnoses in each group. In the EDOU group, the leading diagnoses were: heart failure, pneumonia, urinary tract infection, chest pain, other

specified respiratory disorders, atrial fibrillation and flutter, syncope and collapse, acute exacerbation of chronic obstructive pulmonary disease, unspecified lower respiratory infection, and cerebral infarction.

Multivariable logistic regression analysis, which included the 29 variables that differed significantly between groups, identified five independent predictors of EDOU admission: age ≥80 years, tachypnoea, bradycardia, GCS <15, and anaemia (Table 4, Fig. 2).

Regarding the outcomes, the need for hospital admission was 54% in the EDOU group compared to 15% in the non-EDOU group (aOR = 2.4; 95% CI: 2.0–2.8). The EDOU group also showed higher rates of 30-day rehospitalization (aOR = 1.6; 95% CI: 1.2–2.3). Notably, EDOU care showed a negative, but non-significant, association with 30-day ED

TABLE 3. The 10 most frequent primary emergency diagnoses in the EDOU and no-EDOU groups.

Diagnoses (ICD-10 code)	EDOU N = 6393 n (%)	Diagnoses (ICD-10 code)	Non-EDOU N = 17,562 n (%)
1. Heart failure (I50)	477 (7.5)	1. Back Pain (M54)	697 (4.0)
2. Pneumonia, unspecified (J18.9)	228 (3.6)	2. Abdominal and pelvic pain (R10)	463 (2.6)
3. Urinary tract infection, site not specified (N39.0)	203 (3.2)	3. Urinary tract infection, site not specified (N39.0)	362 (2.1)
4. Chest pain, unspecified (R07.4)	183 (3.0)	4. Heart failure (I50)	358 (2.0)
5. Other specified respiratory disorders (J98.8)	181 (2.8)	5. Injury of unspecified body region (T14)	340 (1.9)
6. Atrial fibrillation and atrial flutter, unspecified (I48.9)	179 (2.8)	6. Other articulation disorders, not elsewhere classified (M25)	280 (1.6)
7. Syncope and collapse (R.55)	173 (2.7)	7. Other specified respiratory disorders (J98)	273 (1.6)
8. Chronic obstructive pulmonary disease with acute exacerbation, unspecified (J44)	158 (2.5)	8. Syncope and collapse (R.55)	239 (1.4)
9. Unspecified acute lower respiratory infection (J22)	155 (2.4)	9. Conjunctivitis (H10)	232 (1.3)
10. Cerebral infarction (I63)	98 (1.5)	10. Airway haemorrhage (haemoptysis, epistaxis) (R04)	211 (1.2)

ICD-10: 10th revision of the International Statistical Classification of Diseases and Related Health Problems; EDOU: Emergency Department Observation Units.

TABLE 4. Results of multivariable logistic regression analysis of the key variables associated with emergency department observation unit admission.

Variables	B	Standard error	Wald	p-value	Adjusted OR	95% CI
Glasgow coma scale score <15 points	1.7	0.4	20.8	<0.001	5.0	2.6–10.0
Significant bradycardia (<50 beats per minute)	1.2	0.5	5.4	0.025	3.1	1.2–8.0
Significant tachypnea (>20 breaths per minute)	0.9	0.2	16.0	<0.001	2.5	1.6–4.0
Anemia (hemoglobin <9 g/dL)	0.7	0.3	6.3	0.014	2.0	1.2–3.6
Age ≥80 yr	0.5	0.1	12.6	<0.001	1.6	1.2–2.1

p-values in bold denote statistical significance ($p < 0.05$).

B: β coefficient; OR: odds ratio; CI: confidence intervals.

revisit among discharged patients (aOR = 0.9; 95% CI: 0.7–1.2).

No significant associations were observed between EDOU use and prolonged hospital stay (aOR = 1.2; 95% CI: 0.9–1.5), 30-day hospital admission after ED discharge (aOR = 1.1; 95% CI: 0.9–1.4), in-hospital mortality (aOR = 1.0; 95% CI: 0.7–1.5), or all-cause 30-day mortality (aOR = 1.0; 95% CI: 0.8–1.4) (Table 5).

4. Discussion

This exploratory study reveals previously unreported sociodemographic, functional, and clinical characteristics of a representative sample of EDOUs in Spain. It also identifies factors associated with admission to EDOUs—an area scarcely addressed in the literature—and describes selected short-term outcomes following observation unit care.

ED visits by older adults are associated with higher rates

of hospital admission, prolonged inpatient stays, unplanned readmissions, functional decline, and mortality [21, 25]. For many of these patients, EDOUs may offer a safe and efficient alternative to conventional admission, helping to avoid unnecessary hospitalization [18].

In our study, patients admitted to EDOUs were older, with greater comorbidity, functional dependence, and cognitive impairment than those who were not admitted to observation units. This suggests a more complex clinical profile, consistent with previous findings [15, 19].

Except for arterial hypertension, all other abnormal physical examination findings were significantly associated with the decision to admit to the EDOU (Table 4). These signs likely influenced clinicians during the triage and initial evaluation phases. Caterino *et al.* [26] found that among older patients in EDOUs, age >65 was not predictive of hospital admission and that hypertension was the only significantly associated vital sign. However, to our knowledge, no previous studies have




FIGURE 2. Adjusted associations for variables analyzed in the EDEN-20 Study with admission in emergency department-observation units (EDOUs). The multivariable model included only significant variables ($p < 0.05$) on univariate analysis. The odds ratio in bold numbers denotes statistically significant differences ($p < 0.05$). ED: Emergency Department; EDOU: Emergency Department Observation Unit; CI: Confidence Interval.

TABLE 5. Association between EDOU admission and short-term outcomes.

Short-term outcomes	EDOUs stay N = 6393 n (%)	Non-EDOU stay N = 17,562 n (%)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	p-value
Need for hospitalization (all patients)	3442 (54.0)	2642 (15.0)	6.6 (6.2–7.0)	2.4 (2.0–2.8)	<0.001
Prolonged hospitalization (>7 d) (in hospitalized patients)	1432 (42.0)	1110 (42.2)	1.0 (0.9–1.1)	1.2 (0.9–1.5)	0.12
30-day revisit ED (in non-hospitalized patients)	706 (24.0)	3083 (21.0)	1.2 (1.0–1.3)	0.9 (0.7–1.2)	0.58
30-day hospital readmission (in hospitalized patients)	385 (11.3)	284 (10.8)	1.0 (0.9–1.2)	1.6 (1.2–2.3)	0.02
30-day hospital admission after discharge from ED (in non-hospitalized patients)	313 (10.6)	864 (5.8)	2.0 (1.7–2.2)	1.1 (0.9–1.4)	0.40
In-hospital mortality (in hospitalized patients)	376 (2.9)	287 (2.9)	1.0 (0.9–1.2)	1.0 (0.7–1.5)	0.80
30-day all-cause mortality (all patients)	590 (9.2)	534 (3.0)	3.2 (2.9–3.7)	1.0 (0.8–1.4)	0.90

Covariates included in the adjusted model were those shown in Table 4.

OR: odds ratio; CI: confidence intervals; ED: Emergency Department; EDOU: Emergency Department Observation Unit. p-values in bold denote statistical significance ($p < 0.05$).

examined the role of physical signs in predicting admission to an observation unit.

Regarding laboratory values, most abnormalities were associated with admission to an EDOU, except for leukopenia, thrombocytopenia, hypoglycaemia, and elevated lactate. Anaemia and erythrocytosis showed the strongest associations, possibly reflecting transfusion needs and exacerbated chronic obstructive pulmonary disease (COPD), respectively.

The analysis of emergency diagnoses (Table 3) showed that the five most common reasons for EDOU admission were heart failure, pneumonia, urinary tract infection, chest pain, and other specified respiratory disorders. Ross *et al.* [15] listed the 20 most common diagnoses among patients aged ≥ 65 admitted to EDOUs in the United States, including conditions such as dehydration, syncope, COPD, cellulitis, abdominal pain, atrial fibrillation, anaemia requiring transfusion, and social problems. All 10 most frequent diagnoses in our study are included within Ross's top 20. Surprisingly, diagnoses like back pain, abdominal/pelvic pain, or trauma-related injuries were not prevalent in our cohort, possibly because these cases are often managed in standard ED consultation areas or referred directly to specialist care.

Our multivariable regression model identified five independent factors associated with EDOU admission: age ≥ 80 years, tachypnoea, bradycardia, GCS < 15 , and anaemia. Although not predictive, the model provides an explanatory framework to identify the clinical profile associated with the decision to admit to EDOUs. Understanding these patterns is key to standardizing practice and optimizing care for older adults in the ED.

Observation unit use among adult ED patients is estimated at around 10% overall [6], and approximately 2% for geriatric patients [15]. In our study, 27% of patients aged ≥ 65 were managed in an EDOU—a proportion rarely reported in the biomedical literature. Furthermore, 54% of hospital admissions occurred after an EDOU stay, compared with just 15% directly from the ED ($p < 0.001$). These figures highlight the central role of EDOUs in managing geriatric patients in Spain.

Previous studies in the U.S. and U.K. have reported inpatient admission rates from EDOUs among older adults ranging from 16% to 78% [15, 27–30]. Our results fall within this range. In unadjusted analyses, 30-day hospital readmission after ED discharge (10.6% vs. 5.8%; OR = 2.0; 95% CI: 1.7–2.2) and 30-day all-cause mortality (9.2% vs. 3.0%; OR = 3.2; 95% CI: 2.9–3.7) were significantly higher in the EDOU group. However, these differences disappeared after adjusting for covariates in the regression model, except for a slight increase in 30-day rehospitalization after observation unit stay (11.3% vs. 10.8%; OR = 1.6; 95% CI: 1.2–1.3; $p < 0.05$).

In an extensive cohort study, Dharmarajan *et al.* [31] evaluated outcomes after observation stays in older adults and reported that 20% of hospital revisits and 50% of all subsequent hospitalizations occurred after an initial observation stay. The 30-day mortality rate in their study was 1.8%, much lower than ours, likely due to the younger age and lower comorbidity burden in their population.

More recently, Berger *et al.* [32] analysed ED return rates after EDOU discharge in a general population and found an overall revisit rate of 9.4%. In contrast, our cohort had a

revisit rate of 24%, possibly reflecting the older age and higher complexity of our patients.

It is important to note that EDOU admission criteria were not standardized across centres. Variability in local protocols, unit structure, and bed availability likely influenced patient selection [33–35]. Therefore, the observed association between EDOU care and outcomes may partly reflect differences in patient complexity rather than the effect of EDOU management itself.

5. Limitations

Our findings should be interpreted in light of several limitations. First, the 48 participating EDOUs were not randomly selected; they volunteered to participate. Nevertheless, they represent 12 of Spain's 17 autonomous communities, including university, high-complexity, and regional hospitals, minimizing potential selection bias.

Second, this analysis is based on a secondary use of a multipurpose registry. Therefore, some associations may be influenced by variables not captured in the original design.

Third, we did not model potential interactions between clinical variables (*e.g.*, bradycardia and beta-blocker use, GCS, and baseline cognitive impairment). Similarly, due to data limitations, we could not apply frailty indices (*e.g.*, Clinical Frailty Scale) or comorbidity scores (*e.g.*, Charlson Index). Instead, we used the Barthel Index as a validated proxy for baseline functional status.

Fourth, we could not confirm whether any individual appeared more than once in the dataset. Although the inclusion period was short, the possibility of repeat visits cannot be excluded. We analysed outcomes per visit, and the data were extracted from electronic health records without patient identifiers.

Lastly, although we described the structural features of each EDOU (Table 1), we did not analyse whether unit type (*e.g.*, Type I vs. Type II) or organizational model (*e.g.*, open vs. closed) influenced outcomes. Future analyses using multilevel or hierarchical models are warranted.

This was an exploratory multicentre registry analysis, and we did not apply formal corrections for multiple comparisons. Although multivariable models were used to control for confounding, univariate associations should be interpreted cautiously. Future studies should confirm these findings in prospectively designed cohorts.

6. Conclusions

Observation units are increasingly used to manage older adults presenting to emergency departments. In this multicentre study involving 48 Spanish hospitals, we found that admission to an EDOU was associated with indicators of greater clinical complexity, such as advanced age, abnormal vital signs, impaired consciousness, and anaemia.

Short-term outcomes, including hospital admission and 30-day readmission, were more frequent among EDOU patients. However, these associations likely reflect patient complexity rather than adverse effects of EDOU care itself. EDOU use was not associated with increased mortality or prolonged hospital

stays.

These findings underscore the need to establish standardized criteria for EDOU admission in geriatric emergency care. Developing evidence-based, protocol-driven strategies may improve patient selection and promote safer, more efficient use of observation resources.

AVAILABILITY OF DATA AND MATERIALS

The data presented in this study are available on reasonable request from the corresponding author.

AUTHOR CONTRIBUTIONS

FJMP—designed the research study. FJMP, IBF, JGC, GBP, ÒM, CF, SA, AAA, JJ, PLL, OJTU, PEH, PPE, MCSP, MRG, MFPP, MGT, MBG, MVB, FLL, RMM, VTG, RJGH, JPG, CLA, MDP, MER, MIV, MRR, AMG—performed the research. FJMP, IBF, JGC, GBP, SA, ÒM—analyzed the data. FJMP, IBF, JGC, ÒM, SA—wrote the manuscript. All authors contributed substantially to the study conception, data acquisition, and data verification, critically revised the manuscript drafts, and read and approved the final version.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The EDEN project received approval from the Clinical Research Ethics Committee of Hospital Clínico San Carlos de Madrid (protocol HCSC/22/005-E). Given the registry's non-interventional nature, Spanish legislation allows centralized approval by a lead ethics committee, with required notification to local committees. The requirement for informed consent was waived by the same Ethics Committee due to the study's retrospective, non-interventional design and the use of anonymized data. The study adhered to the principles outlined in the Declaration of Helsinki.

ACKNOWLEDGMENT

We thank all members of the SIESTA network for their commitment and valuable contributions to the development of this study.

SIESTA-Network Researchers:

Hospital Clínico San Carlos, Madrid: Juan González del Castillo, Cesáreo Fernández Alonso, Jorge García Lamberechts, Alejandro Melcon Villalibre, Sara Vargas Lobé, Laura Fernández García, Beatriz Escudero Blázquez, Estrella Serrano Molina, Julia Barrado Cuchillo.

Hospital Universitario Infanta Cristina, Parla: Ángel Iván Díaz Salado, Alicia Fuente Gaforio, Cristina Güemes de la Iglesia, Beatriz Honrado Galán.

Hospital Santa Tecla, Tarragona: Enrique Martín Mojarro, Sílvia Flores Quesada, Lidia Cuevas Jiménez.

Hospital Universitario de Canarias, Tenerife: Guillermo Burillo Putze, Aarati Vaswani-Bulchand, Patricia Eiroa Hernández.

Hospital Norte Tenerife: Patricia Parra-Esquivel, Montserrat Rodríguez-Cabrera.

Hospital General Universitario Reina Sofía, Murcia: Pascual Piñera Salmerón, Lorena Bernabé Vera, Juan José López Pérez, Paula Lázaro Aragüés, Mariya Ovsepyan, M. Encarnación Sánchez Cánovas.

Hospital Universitario del Henares, Madrid: David Ampuero Martinich, Martín Ruiz Grinspan, Cristóbal Rodríguez Leal, Rocío Martínez Avilés.

Hospital Clínic, Barcelona: Òscar Mir, Sònia Jiménez, Sira Aguiló Mir, Francesc Xavier Alemany González, María Florencia Poblete Palacios, Claudia Lorena Amarilla Molinas, Ivet Gina Osorio Quispe, Sandra Cuerpo Cardeñosa.

Hospital General Universitario de Elche: Matilde González Tejera, Ana Puche Alcaraz, Cristina Chacón García.

Hospital Universitario y Politécnico La Fe, Valencia: Leticia Serrano Lázaro, Javier Millán Soria, Jérica Mansilla Collado, María Bóveda García.

Hospital Universitario Dr Balmis, Alicante: Pere Llorens Soriano, Adriana Gil Rodrigo, Begoña Espinosa Fernández, Mónica Veguillas Benito, Sergio Guzmán Martínez, Gema Jara Torres, María Caballero Martínez.

Hospital Universitario de Bellvitge, L'Hospitalet de Llobregat, Barcelona: Javier Jacob Rodríguez, Ferran Llopis, Elena Fuentes, Lidia Fuentes, Francisco Chamorro, Lara Guillen, Nieves López.

Hospital de Axarquía, Málaga: Coral Suero Méndez, Lucía Zambrano Serrano, Rocío Muñoz Martos, Rocío Lorenzo Álvarez.

Hospital Regional Universitario de Málaga: Manuel Salido Mota, Valle Toro Gallardo, Antonio Real López, Lucía Ocaña Martínez, Esther Muñoz Soler, Mario Lozano Sánchez, Eva María Fragero Blesa.

Hospital Santa Barbara, Soria: Fahd Beddar Chaib, Rodrigo Javier Gil Hernández.

Hospital Valle de los Pedroches, Pozoblanco, Córdoba: Jorge Pedraza García, Paula Pedraza Ramírez.

Hospital Universitario Reina Sofía, Córdoba: F. Javier Montero-Pérez, Carmen Lucena Aguilera, F. de Borja Quero Espinosa, Ángela María Cobos Requena, Esperanza Muñoz Triano, Inmaculada Bajo Fernández, María Calderón Caro, Sierra Bretones Baena.

Hospital Universitario Gregorio Marañón, Madrid: Juan Fernández Herranz, Marta Rincón Francés, Irene Arnaiz Fernández Esther Gargallo García, Juan Antonio Andueza Lillo, Iria Miguens Blanco, Ioana Muñoz Betegón, Dariela Edith Micheloud Giménez.

Hospital Universitario de Burgos: Verónica Castro Jiménez, Lucía González Ferreira, Rocio Hernando González, María Eugenia Rodríguez Palma, Javier Varona Castrillo.

Complejo Asistencial Universitario de León: Albert Carbó Jordá, Enrique González Revuelta, Héctor Lago Gancedo, Miguel Moreno Martín, Alberto Álvarez Madrigal, Marta Iglesias Vela, Mónica Santos Orús.

Hospital Universitario Morales Meseguer, Murcia: Rafael Antonio Pérez-Costa, María Rodríguez Romero, Esperanza Marín Arranz, Ana Barnes Parra.

Hospital Francesc de Borja de Gandía, Valencia: Natalia Morera Sendra, Beatriz González García, Fátima Peiró Monzó.

Hospital Universitario Severo Ochoa, Leganés. Madrid: María José Hernández Martínez, Ana Benito Blanco, Vanesa Abad Cuñado, Julia Martínez-Ibarreta Zorita, Irene Cabrera Rodrigo.

Hospital Clínico Universitario Virgen Arrixaca, Murcia: Miguel Parra Morata Eva Quero Motto, Nuria Tomas Garcia, Laura Bernal Martínez.

Hospital Universitario Lorenzo Guirao, Cieza, Murcia: Alberto Artieda Larrañaga, José Joaquín Giménez Belló.

Hospital Universitario Dr. Josep Trueta, Girona: María Adroher Muñoz, Ester Soy Ferrer, Eduard Anton Poch Ferrer.

Hospital de Mendaro, Guipuzkoa: Jeong-Uh Hong Cho.

Hospital Universitario Miguel Servet, Zaragoza: Patricia Trenc Español, Fernando López López, Jorge Navarro Calzada, Belén Gros Bañeres, Cristina Martín Durán, María Teresa Escolar Martínez-Berganza, Iciar González Salvatierra.

Hospital Comarcal El Escorial, Madrid: Sara Gayoso Martín.

Hospital Do Salnes, Villagarcía de Arosa, Pontevedra: María Goretti Sánchez Sindín.

Hospital de Barbanza, Ribeira, A Coruña: Martina Silva Penas, Azucena Prieto Zapico.

Hospital del Mar, Barcelona: Bárbara Gómez y Gómez, Isabel Cirera Lorenzo, Patricia Gallardo Vizcaino, Margarita Puigali Ballard.

Hospital Santa Creu y Sant Pau, Barcelona: Aitor Alquezar Arbé, Carlos Romero Carrete, Sergio Pérez Baena, Laura Lozano Polo, Roser Arenos Sambro, José María Guardiola Tey, Carme Beltrán Vilagrasa.

Hospital de Vic, Barcelona: Lluís Llauger.

Hospital Valle del Nalón, Langreo, Asturias: Ángel Peláez González, Celia Rodríguez Valles.

Hospital Altagracia, Manzanares, Ciudad Real: Laura Márquez Quero.

Hospital Nuestra Señora del Prado de Talavera de la Reina, Toledo: Laura Molina, Mónica Cañete, Ricardo Juárez González.

Hospital Universitario Vinalopó, Elche, Alicante: María Martínez Juan María José Blanco Hoffman, Pedro Ruiz Asensio.

Hospital de Móstoles, Madrid: Fátima Fernández Salgado, Eva de las Nieves Rodríguez, Gema Gómez García.

Hospital Virgen del Rocío, Sevilla: Rafaela Ríos Gallardo, Teresa Pablos Pizarro, Mariano Herranz García, Laura Redondo Lora, Ana Gómez Caminero, Claudio Bueno Mariscal, Amparo Fernández-Simón Almela, Esther Pérez García.

Hospital General Universitario Dr. Peset. Valencia. María Amparo Berenguer Diez. María Ángeles de Juan Gómez. María Luisa López Grima, Rigoberto Jesús del Río Navarro.

Hospital Universitario Son Espases. Palma de Mallorca: Núria Perelló Viola. Bernardino Comas Diaz, Sandra Guiu Martí, Juan Domínguez Casasola.

Clínica Universitaria Navarra, Madrid: Nieves López-Laguna.

Hospital Clínico Universitario de Valencia: José J. Noceda Bermejo, María Teresa Sánchez Moreno, Raquel Benavent Campos, Jacinto García Acosta, Alejandro Cortés Soler.

Hospital Alvaro Cunqueiro, Vigo: María Teresa Maza Vera, Raquel Rodríguez Calveiro, Paz Balado Dacosta, Violeta Del-

gado Sardina, Emma González Nespereira, Carmen Fernández Domato, Elena Sánchez Fernández-Linares.

Hospital Universitario de Salamanca: Ángel García García, Francisco Javier Diego Robledo, Manuel Ángel Palomero Martín, Jesús Ángel Sánchez Serrano.

Hospital de Zumarraga, Gipuzkoa: Patxi Ezponda.

Hospital Virxe da Xunqueira, La Coruña: Andrea Martínez Lorenzo.

Hospital Universitario Río Hortega, Valladolid: Inmaculada García Rupérez, Pablo González Garcinuño, Raquel Hernando Fernández, José Ramón Oliva Ramos, Virginia Carbajosa Rodríguez.

Hospital Central Asturias: Alba Martínez Alonso, Angela López Carrillo, Belén Pérez Fernández, Carmen Pérez Fonsaca, Claudia Corugedo Ovies, Claudia Marinero Noval, Eugenia Prieto Piquero.

Hospital Juan Ramón Jiménez, Huelva: Asumpta Ruiz Aranda, María José Marchena, José María Santos Martín.

FUNDING

This research received no external funding.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- [1] Magidson PD, Carpenter CR. Trends in geriatric emergency medicine. *Emergency Medicine Clinics of North America*. 2021; 39: 243–255.
- [2] Quality improvement initiatives for the care of geriatric patients in the emergency department. *Annals of Emergency Medicine*. 2016; 68: 404.
- [3] Miró Ò, González Del Castillo J. Collaboration among Spanish emergency departments to promote research: on the creation of the SIESTA (Spanish Investigators in Emergency Situations TeAm) network and the coordination of the UMC-19 (Unusual Manifestations of COVID-19) macropoint. *Emergencias*. 2020; 32: 269–277.
- [4] Miró Ò, González Del Castillo J. SIESTA—the Spanish Investigators in Emergency Situations TeAm—during the COVID-19 pandemic: a summary of results of the network's first challenge. *Emergencias*. 2022; 34: 225–227.
- [5] Kahn JH, Magauran BG, Olshaker JS, Shankar KN. Current trends in geriatric emergency medicine. *Emergency Medicine Clinics of North America*. 2016; 34: 435–452.
- [6] Kabell Nissen S, Rueegg M, Carpenter CR, Kaeppli T, Busch J, Fournaise A, *et al.* Prognosis for older people at presentation to emergency department based on frailty and aggregated vital signs. *Journal of the American Geriatrics Society*. 2023; 71: 1250–1258.
- [7] Hustey FM. Geriatric observation medicine. In Mace SE (ed.) *Observation medicine: principles and protocols* (pp. 304–308). 1st edn. Cambridge University Press: Cambridge, UK. 2017.
- [8] Miró Ò, Morales X, Cuerpo S, Möckel M, Burillo G, Alquézar-Arbé A, *et al.*; SIESTA research network members. Non-specific abdominal pain in elderly patients discharged from the emergency department: frequency, outcomes and risk-factors for adverse events (EDEN-43 study). *The American Journal of Emergency Medicine*. 2025; 93: 140–145.
- [9] Rider I, Sorensen M, Brady WJ, Gottlieb M, Benson S, Koyfman A, *et al.* Disposition of acute decompensated heart failure from the emergency department: an evidence-based review. *The American Journal of Emergency Medicine*. 2021; 50: 459–465.
- [10] Saviano A, Zanza C, Longhitano Y, Ojetti V, Franceschi F, Bellou A, *et al.* Current trends for delirium screening within the emergency department. *Medicina*. 2023; 59: 1634.

- [11] Gruenberg CC, Breaud AH, Liu JH, Mitchell PM, Feldman JA, Nelson KP, *et al.* Are geriatric patients placed in an emergency department observation unit on a chest pain pathway more likely than non-geriatric patients to re-present to the hospital within 30 days? *The Journal of Emergency Medicine*. 2018; 54: 302–306.
- [12] Navas A, Guzman B, Hassan A, Borawski JB, Harrison D, Manandhar P, *et al.* Untapped potential for emergency department observation unit use: a national hospital ambulatory medical care survey (NHAMCS) study. *The Western Journal of Emergency Medicine*. 2022; 23: 134–140.
- [13] Gabayan GZ, Liang LJ, Doyle B, Huang DY, Sarkisian CA. Emergency department increased use of observation care for elderly Medicare patients. *Journal of Hospital Administration*. 2018; 7: 9–16.
- [14] Perry M, Franks N, Pitts SR, Moran TP, Osborne A, Peterson D, *et al.* The impact of emergency department observation units on a health system. *The American Journal of Emergency Medicine*. 2021; 48: 231–237.
- [15] Ross MA, Compton S, Richardson D, Jones R, Nittis T, Wilson A. The use and effectiveness of an emergency department observation unit for elderly patients. *Annals of Emergency Medicine*. 2003; 41: 668–677.
- [16] Nouvenne A, Ticinesi A, Cerundolo N, Prati B, Parise A, Chiussi G, *et al.* Implementing a multidisciplinary rapid geriatric observation unit for non-critical older patients referred to hospital: observational study on real-world data. *Aging Clinical and Experimental Research*. 2022; 34: 599–609.
- [17] Roberts MV, Baird W, Kerr P, O'Reilly S. Can an emergency department-based clinical decision unit successfully utilize alternatives to emergency hospitalization? *European Journal of Emergency Medicine*. 2010; 17: 89–96.
- [18] Blecker S, Gavin NP, Park H, Ladapo JA, Katz SD. Observation units as substitutes for hospitalization or home discharge. *Annals of Emergency Medicine*. 2016; 67: 706–713.e2.
- [19] Bell J, Lim S, Mikami T, Bahk J, Argiro S, Steiger D. The impact on thirty day readmissions for patients hospitalized for acute exacerbations of chronic obstructive pulmonary disease admitted to an observation unit versus an inpatient medical unit: a retrospective observational study. *Chronic Respiratory Disease*. 2024; 21: 14799731241242490.
- [20] Aminzadeh F, Dalziel WB. Older adults in the emergency department: a systematic review of patterns of use, adverse outcomes, and effectiveness of interventions. *Annals of Emergency Medicine*. 2002; 39: 238–247.
- [21] Curiati PK, Gil-Junior LA, Morinaga CV, Ganem F, Curiati JAE, Avelino-Silva TJ. Predicting hospital admission and prolonged length of stay in older adults in the emergency department: the PRO-AGE scoring system. *Annals of Emergency Medicine*. 2020; 76: 255–265.
- [22] Miró Ò, Jacob J, García-Lamberechts EJ, Piñera Salmerón P, Llorens P, Jiménez S, *et al.*; Red de investigación SIESTA. Sociodemographic characteristics, functional status, and health resource use of older patients treated in Spanish emergency departments: a description of the EDEN cohort. *Emergencias*. 2022; 34: 418–427.
- [23] Montero-Pérez FJ, Cobos Requena ÁM, González Del Castillo J, Jacob J, García-Lamberechts EJ, Piñera Salmerón P, *et al.*; Investigadores de la red SIESTA. Impact of the COVID-19 pandemic on demand for emergency department care for older patients: the EDEN-7 COVID cohort study. *Emergencias*. 2023; 35: 270–278.
- [24] Martín-Sánchez FJ, Bermejo Boixareu C. EDEN—the Emergency Department and Elder Needs project—is a chance to understand and improve a whole-system approach to elder care in Spanish emergency departments. *Emergencias*. 2022; 34: 409–410.
- [25] Hominick K, McLeod V, Rockwood K. Characteristics of older adults admitted to hospital versus those discharged home, in emergency department patients referred to internal medicine. *Canadian Geriatrics Journal*. 2016; 19: 9–14.
- [26] Caterino JM, Hoover EM, Moseley MG. Effect of advanced age and vital signs on admission from an ED observation unit. *The American Journal of Emergency Medicine*. 2013; 31: 1–7.
- [27] Harrop SN, Morgan WJ. Emergency care of the elderly in the short-stay ward of the accident and emergency department. *Emergency Medicine Journal*. 1985; 2: 141–147.
- [28] Khan SA, Millington H, Miskelly FG. Benefits of an accident and emergency short stay ward in the staged hospital care of elderly patients. *Emergency Medicine Journal*. 1997; 14: 151–152.
- [29] Madsen TE, Bledsoe J, Bossart P. Appropriately screened geriatric chest pain in an observation unit are not admitted at a higher rate than nongeriatric patients. *Critical Pathways in Cardiology*. 2008; 7: 245–247.
- [30] Mace SE, Graff L, Mikhail M, Ross M. A national survey of observation units in the United States. *The American Journal of Emergency Medicine*. 2003; 21: 529–533.
- [31] Dharmarajan K, Qin L, Bierlein M, Choi JES, Lin Z, Desai NR, *et al.* Outcomes after observation stays among older adult Medicare beneficiaries in the USA: retrospective cohort study. *The BMJ*. 2017; 357: j2616.
- [32] Berger D, King S, Caldwell C, Soto EF, Chambers A, Boehmer S, *et al.* Returns after discharge from the emergency department observation unit: who, what, when, and why? *The Western Journal of Emergency Medicine*. 2023; 24: 390–395.
- [33] Aguiló Mir S. Identifying the frail patient in the emergency department: an urgent and necessary effort. *Emergencias*. 2023; 35: 165–166.
- [34] Fernández Alonso C, Del Arco Galán C, Torres Garate R, Madrigal Valdés JF, Romero Pareja R, Bibiano Guillén C, *et al.*; Registro Frail-ED-Madrid. Performance of 3 frailty scales for predicting adverse outcomes at 30 days in older patients discharged from emergency departments. *Emergencias*. 2023; 35: 196–204.
- [35] Puig-Campmany M, Ris Romeu J. Frail older patients in the emergency department: main challenges. *Emergencias*. 2022; 34: 415–417.

How to cite this article: Francisco Javier Montero-Pérez, Inmaculada Bajo-Fernández, Juan González del Castillo, Guillermo Burillo-Putze, Cesáreo Fernández, Sira Aguiló, *et al.*, on behalf of the researchers of the SIESTA network. Admission rates and outcomes of elderly patients in emergency department observation units. *A Spanish multicentre study*. *Signa Vitae*. 2025; 21(10): 80-91. doi: 10.22514/sv.2025.146.