




CORRECTION

Correction: Comparison of modified thoracoabdominal nerve block through perichondral approach (M-TAPA block) and trocar site local anesthetic infiltration for pain management in total laparoscopic hysterectomy: a randomized-controlled trial

Ela Erten¹, Umut Kara^{2,*}, Fatih Simsek¹, Mehmet Emin İnce¹,
Vildan Altahhan³, Ozhan Ozdemir³, Gokhan Berktug Bahadır⁴

¹Department of Anesthesiology and Reanimation, Gulhane Training and Research Hospital, 06010 Ankara, Turkey

²Department of Anesthesiology and Reanimation, Acıbadem Hospital Atasehir, 34758 Istanbul, Turkey

³Department of Gynaecology and Obstetrics, Gulhane Training and Research Hospital, 06010 Ankara, Turkey

⁴Department of Paediatric Surgery, Gulhane Training and Research Hospital, 06010 Ankara, Turkey

*Correspondence

drumutkara@gmail.com (Umut Kara)

Error in author name

The authors regret that the fifth author's name was incorrectly written as Vildan Atasever in the original publication [1]. The correct name should read as Vildan Altahhan. The authors would like to apologise for any inconvenience caused.

REFERENCES

- [1] Erten E, Kara U, Simsek F, İnce ME, Altahhan V, Ozdemir O, *et al.* Comparison of modified thoracoabdominal nerve block through

perichondral approach (M-TAPA block) and trocar site local anesthetic infiltration for pain management in total laparoscopic hysterectomy: a randomized-controlled trial. *Signa Vitae*. 2025; 21: 99–106.

How to cite this article: Ela Erten, Umut Kara, Fatih Simsek, Mehmet Emin İnce, Vildan Altahhan, Ozhan Ozdemir, *et al.* Correction: Comparison of modified thoracoabdominal nerve block through perichondral approach (M-TAPA block) and trocar site local anesthetic infiltration for pain management in total laparoscopic hysterectomy: a randomized-controlled trial. *Signa Vitae*. 2026; 22(1): 147-147. doi: 10.22514/sv.2026.007.