

CORRECTION



Correction: Guidelines-recommended tidal volumes are not achieved during continuous mechanical chest compressions—results from a laboratory study

Matej Mažič^{1,2,*}, Andrej Markota^{2,3}

¹Department of Anesthesiology and Perioperative Intensive Medicine, General and Teaching Hospital Celje, 3000 Celje, Slovenia

²Faculty of Health Sciences, University of Novo mesto, 8000 Novo mesto, Slovenia

³Department of Internal Intensive Medicine, University Medical Center Maribor, 2000 Maribor, Slovenia

*Correspondence

matej.mazic@sb-celje.si (Matej Mažič)

In the originally published article [1] (DOI: 10.22514/sv.2025.030), the authors' affiliations were incomplete.

The affiliations have been corrected as follows:

Matej Mažič^{1,2,*}, Andrej Markota^{2,3}

¹Department of Anesthesiology and Perioperative Intensive Medicine, General and Teaching Hospital Celje, 3000 Celje, Slovenia

²Faculty of Health Sciences, University of Novo mesto, 8000 Novo mesto, Slovenia

³Department of Internal Intensive Medicine, University Medical Center Maribor, 2000 Maribor, Slovenia

*Corresponding Author.

The authors confirm that this correction does not affect the scientific content, results, or conclusions of the study.

This Correction has been approved by the Editor-in-Chief. The original article PDF remains available as published and has not been altered. Readers are advised to refer to this Correction for the updated author affiliations.

REFERENCES

- [1] Mažič M, Markota A. Guidelines-recommended tidal volumes are not achieved during continuous mechanical chest compressions—results from a laboratory study. *Signa Vitae*. 2025; 21: 1–7.